arXiv:0905.1741v1 [math.AG] 12 May 2009

ON THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF LINEAR TORUS CURVES OF MAXIMAL CONTACT

M. KAWASHIMA

Abstract. In this paper, we compute the fundamental group of the complement of linear torus curves of maximal contact and we show that it is isomorphic that of generic linear torus curves.

1. INTRODUCTION

A plane curve $C \subset \mathbb{P}^2$ of degree pq is called a *curve of* (p,q) *torus type* with $p > q \geq 2$, if there is a defining polynomial F of C of the form $F = F_p^q - F_q^p$, where F_p, F_q are homogeneous polynomials of X, Y, Z of degree p and q respectively. A singular point $P \in C$ is called *inner* if $P \in \{F_p = F_q = 0\}$. Otherwise, P is called an *outer singularity.* C is called *tame* if C has no outer singularity.

We say that a (p,q) torus curve C is *a torus curve of a maximal contact* if $\{F_p = F_q = 0\} = \{\xi_0\}$ and $\{F_p = 0\}$ is smooth at ξ_0. Then the intersection multiplicity $I(F_q, F_p; \xi_0)$ is equal to pq by Bézout's theorem and singularity (C, ξ_0) is topologically equivalent to the Brieskorn-Pham singularity $B_{p^2q,q}$ where $B_{m,n} := \{(x,y) \in \mathbb{C}^2 \mid x^m + y^n = 0\}$.

We are interested in topological invariants the Alexander polynomial $\Delta_C(t)$ and the fundamental groups $\pi_1(\mathbb{P}^2 \setminus C)$ and $\pi_1(\mathbb{C}^2 \setminus C)$. It is known that if C is a tame maximal contact of type (p,q), then the Alexander polynomial of C is equal to

$$\Delta_C(t) = \frac{(t^{pq/r} - 1)^r (t-1)}{(t^p - 1)(t^q - 1)}$$

where $r = \gcd(p,q)$ ([1, 2]).

We consider the special class of torus curves of type (p,q). We say that C is *a linear torus curve of type* (p,q) if $F_q = L^q$ for some linear form L. If C is a linear torus curve of type (p,q), then C generically consists of q components of curves of degree p as

$$F = \prod_{j=1}^{q} (F_p - \zeta^j L^p), \quad \text{where} \quad \zeta := \exp\left(\frac{2\pi\sqrt{-1}}{q}\right)$$

and the inner singularities of C are contained in the intersection $C_p \cap L$. If $C_p \cap L$ consists of p distinct points, C is called *a generic linear torus curve.* In this case, C has p singular points, each of which is topologically equivalent to $B_{pq,q}$.

If C is a generic linear torus curve of type (p,q), then C has q components of degree p which are defined by $C^{(j)} : \{F_p - \zeta^j L^p = 0\}$ for $j = 1, \dots, q$. Then $C^{(i)}$ and $C^{(j)}$ are tangent at p points $\{F_p = L = 0\}$ with the intersection multiplicity p each other. The fundamental group is given by

$$\pi_1(\mathbb{P}^2 \setminus C) \cong F(q-1) * \mathbb{Z}/p\mathbb{Z}$$

2000 *Mathematics Subject Classification.* 14H20,14H30,14H45.

Key words and phrases. Torus curve, maximal contact.

where $F(d)$ is the free group of rank d ([4]). Its Alexander polynomial $\Delta_C(t)$ is given by

$$\Delta_C(t) = \frac{(t^{pq}-1)^{q-1}(t-1)}{t^q-1}.$$

Let C be a tame (p,q) linear torus curve of a maximal contact. Then C has q components of degree p which intersect at one point with intersection multiplicity p^2 each other. In this paper, we compute that fundamental groups of $\mathbb{P}^2 \setminus C$ and $\mathbb{C}^2 \setminus C$.

Our main result is the following:

Theorem 1. *Let C be a tame (p,q) linear torus curve of a maximal contact. Then the fundamental group $\pi(\mathbb{P}^2 \setminus C)$ is isomorphic to that of generic linear torus curves. Namely*

$$\pi(\mathbb{C}^2 \setminus C) \cong \langle g_1, \dots, g_q, \omega \,|\, \omega = g_1 \cdots g_q, [g_j, \omega^p] = e, j = 1, \dots, q\rangle$$
$$\pi(\mathbb{P}^2 \setminus C) \cong \langle g_1, \dots, g_q, \omega \,|\, \omega^p = e,\ \omega = g_1 \cdots g_q\rangle \cong F(q-1) * \mathbb{Z}/p\mathbb{Z}.$$

2. PRELIMINARIES

2.1. **Van Kampen-Zariski Pencil method.** Let C be a reduced plane curve of degree d in $\mathbb{P}^2$. To compute the fundamental groups $\pi_1(\mathbb{P}^2 \setminus C)$ and $\pi_1(\mathbb{C}^2 \setminus C)$, we use so-called *Van Kampen-Zariski pencil method.* We recall it briefly in the following ([6]). We fix a point $B_0 \in \mathbb{P}^2 \setminus C$ and we consider the set of lines $\mathcal{L} = \{L_s \,|\, s \in \mathbb{P}^1\}$ through B_0 and $\mathcal{L}$ is called a *pencil.* Taking a linear change of coordinates if necessary, we may assume that $B_0 = [1:0:0]$ and L_s is defined by $L_s = \{Y - sZ = 0\}$ in $\mathbb{P}^2$ where (X, Y, Z) is the fixed homogeneous coordinates. Take $L_\infty = \{Z = 0\}$ as the line at infinity and assume that L_∞ intersects transversely with C. We consider the affine coordinates $(x, y) = (X/Z, Y/Z)$ on $\mathbb{C}^2 = \mathbb{P}^2 - L_\infty$. Let $F(X, Y, Z)$ be the defining homogeneous polynomial of C and let $f(x, y) = F(x, y, 1)$ be the affine equation of C. We use the following notations:

$$C^a = C \cap \mathbb{C}^2, \quad L_s^a = L_s \cap \mathbb{C}^2.$$

We identify L_s and L_s^a with $\mathbb{P}^1$ and $\mathbb{C}$ respectively and the pencil line L_s^a is defined by $\{y = s\}$ in the affine coordinates (x, y). We use x as the coordinates of L_s^a.

A pencil line L_s is called *singular with respect to C* if L_s passes through a singular point of C or L_s is tangent to C. Otherwise, we call L_s is *generic.* Hearafter we assume that L_∞ is generic and B_0 is not contained in C.

Let $\mathbb{C}_y$ be the space of the parameters of the pencil with coordinates y and let $\Sigma = \{s \in \mathbb{C}_y \,|\, L_s \text{ is a singular pencil lines}\}$ and suppose that $\Sigma = \{s_1, \dots, s_k\} \subset \mathbb{C}_y$. We fix a generic pencil line L_{s_0} (so $s_0 \in \mathbb{C}_y \setminus \Sigma$) and put $L_{s_0}^a \cap C^a = \{Q_1, \dots, Q_d\}$ where d is the degree of C. We take a base point $*_0 \in L_{s_0}^a \setminus L_{s_0}^a \cap C^a$ on the real axis which is sufficiently near to B_0 and $*_0 \neq B_0$. We take a large disk $\Delta_R \subset L_{s_0}^a$ such that $L_{s_0}^a \cap C^a \subset \Delta_R$ and $*_0 \notin \Delta_R$. We may assume that $\Delta_R = \{(x, s_0) \in L_{s_0}^a \,|\, |x| \leq R\}$ with a sufficient large R. We orient the boundary of Δ_R counter-clockwise and we put $\Xi = \partial\Delta_R$. Join the circle Ξ to the base point by a line segment L connecting $*_0$ and Ξ along the real axis. Let Ω be the class of this loop $L \circ \Xi \circ L^{-1}$ in $\pi_1(L_{s_0}^a \setminus L_{s_0}^a \cap C; *_0)$. We take free generators $g_1, \dots, g_d$ of $\pi_1(L_{s_0}^a \setminus L_{s_0}^a \cap C; *_0)$ so that g_i goes around Q_i counter-clockwise along a small circle and we assume that $\omega = g_d \cdots g_1$, taking a suitable ordering of $g_1, \dots, g_d$ if necessary.



FIGURE 1. $L_{s_0} \cap C$

Hereafter we denote a small lasso oriented in the counter clockwise direction by a bullet with a path in the following figures. Thus ——● indicates ——↺.

The fundamental group $\pi_1(\mathbb{C}_y \setminus \Sigma; s_0)$ acts on $\pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)$ which we call *the monodromy action* of $\pi_1(\mathbb{C}_y \setminus \Sigma; s_0)$. For details, we refer to [6, 5]. Note that $\pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)$ is a free group of rank d with generators $g_1, \dots, g_d$. The class of the action of $\sigma \in \pi_1(\mathbb{C}_y \setminus \Sigma; s_0)$ on $g \in \pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)$ is denoted by g^σ.

Let $\mathcal{M}$ be the normal subgroup of $\pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)$ which is normally generated by

$$\mathcal{R} = \langle g^{-1} g^\sigma \mid g \in \pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0),\ \sigma \in \pi_1(\mathbb{C}_y \setminus \Sigma; s_0) \rangle$$

and we call $\mathcal{M}$ the *the group of the monodromy relations*. Put

$$\mathcal{M}(\sigma_i) = \{ g_j^{-1} g_j^{\sigma_i} \mid j = 1, \dots, d \}.$$

Then the group $\mathcal{M}$ is the minimal normal subgroup of $\pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)$ generated by $\bigcup_{j=1}^k \mathcal{M}(\sigma_i)$. By the definition, we have the relation

$$R(\sigma_i) \qquad\qquad g_j = g_j^{\sigma_i}$$

in the quotient group $\pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)/\mathcal{M}$. We call $R(\sigma_i)$ *the monodromy relation for* σ_i. Let $j : L^a_{s_0} \setminus L^a_{s_0} \cap C \to \mathbb{C}^2 \setminus C^a$ and $\iota : \mathbb{C}^2 \setminus C^a \to \mathbb{P}^2 \setminus C$ be the respective inclusions.

Proposition 1 ([9, 8, 7])**.** *Under the above situations,*

(1) *The canonical homomorphism* $j_\sharp : \pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0) \to \pi_1(\mathbb{C}^2 \setminus C^a; *_0)$ *is surjective and the kernel* $\mathrm{Ker} j_\sharp$ *is equal to* $\mathcal{M}$. *Thus we have the isomorphism:*

$$\pi_1(\mathbb{C}^2 \setminus C^a; *_0) \cong \pi_1(L^a_{s_0} \setminus L^a_{s_0} \cap C; *_0)/\mathcal{M}.$$

(2) ([3]) *The canonical homomorphism* $\iota_\sharp : \pi_1(\mathbb{C}^2 \setminus C^a; *_0) \to \pi_1(\mathbb{P}^2 \setminus C; *_0)$ *is surjective and the kernel* $\mathrm{Ker}\iota_\sharp$ *is generated by a single element* $\omega = g_d \cdots g_1$ *which is in the center of* $\pi(\mathbb{C}^2 \setminus C^a)$ *and* $\mathrm{Ker}\iota_\sharp = \langle \omega \rangle \cong \mathbb{Z}$. *Thus we have an isomorphism*

$$\pi_1(\mathbb{P}^2 \setminus C; *_0) \cong \pi_1(\mathbb{C}^2 \setminus C^a; *_0)/\langle \omega \rangle$$

3. PROOF OF THEOREM 1

Let (x, y) be affine coordinates such that $x = X/Z$, $y = Y/Z$ on $\mathbb{C}^2 := \mathbb{P}^2 \setminus \{Z = 0\}$.

3.1. **Construction of curves.** In this section, we construct a linear torus curve C of a maximal contact and investigate its local properties. First we introduce a plane curve $D_\alpha = \{g_\alpha(x,y)=0\}$ of degree p where the defining polynomial $g_\alpha(x,y)$ is defined by

$$g_\alpha(u,y) = u - \psi(y,\alpha), \quad \psi(y,\alpha) = y - \alpha y^p, \quad \alpha \in \mathbb{C}^*.$$

Now we consider the p-fold cyclic covering ([5]) which is defined by

$$\varphi_p : \mathbb{C}^2 \to \mathbb{C}^2, \quad \varphi_p(x,y) = (u,y), \quad u = x^p.$$

To distinguish two affine planes, we denote the source space of φ_p by $\mathbb{C}^2_s$ with coordinates (x,y) and the target space of φ_p by $\mathbb{C}^2_t$ with coordinates (u,y). Hereafter we simply denote $\mathbb{C}^2$ instead of $\mathbb{C}^2_s$.

Let $C_\alpha := \varphi^{-1}(D_\alpha)$ be the pull-back of D_α by φ_p and let $f_\alpha(x,y) = g_\alpha(x^p,y)$ be the defining polynomial of C_α. Note that

$$f_\alpha(x,y) = x^p - \psi(y,\alpha).$$

By the defining equation of C_α, we see that the set of parameters which corresponds to the singular pencil lines for C_α are given by

$$\Sigma_\alpha := \{y \in \mathbb{C}_y \mid \psi(y,\alpha)=0\}$$

(cf. [4]). Fix a γ such that $\gamma^{p-1} = 1/\alpha$. Then we factorize $\psi(y,\alpha)$ as follows:

$$\psi(y,\alpha) = \alpha\, y \prod_{k=0}^{p-2}(\gamma\xi^k - y), \quad \xi := \exp\left(\frac{2\pi\sqrt{-1}}{p-1}\right).$$

Then we can see that $O=(0,0)$ and $Q_k = (0,\gamma\,\xi^k)$ for $k=0,\dots,p-2$ are flex points of C_α of flex order $p-2$ and their tangent lines are nothing but the singular pencil lines through these points and they are given by $y=0$ and $y=\gamma\xi^k$ respectively.

Now we are ready to define a reduced curve C. Take q non-zero mutually distinct complex numbers $\alpha_1,\dots,\alpha_q$ and put $D_j = \{g_{\alpha_j}(x,y)=0\}$ for $j=1,\dots,q$ and put $D = \bigcup_{i=1}^q D_j$. Then put $C_j = \varphi_p^{-1}(D_j)$ for $=1,\dots,q$ and finally we define

$$C = \varphi_p^{-1}(D) = C_1 \cup \cdots \cup C_q.$$

The defining polynomials $f_j(x,y)$ and $f(x,y)$ of C_j and C respectively are given as follows.

$$f_j(x,y) = x^p - \psi(y,\alpha_j), \quad f(x,y) = \prod_{j=1}^q f_j(x,y).$$

Put $U = \{(\alpha_1,\dots,\alpha_q) \in \mathbb{C}^{*q} \mid \alpha_i \neq \alpha_j, \text{ for any } i \neq j\}$. It is known that the embedded topology of $C \subset \mathbb{C}^2$ does not depend on the choice of $(\alpha_1,\dots,\alpha_q) \in U$ (see [2]).

Lemma 1. *The reduced curve C can be a linear torus curve of a maximal contact for a certain choice of $(\alpha_1,\dots,\alpha_q)$.*

Proof. We take $(\alpha_1,\dots,\alpha_q) = (1,\zeta,\dots,\zeta^{p-1}) \in U$, then we claim that $C = \{f(x,y)=0\}$ is a (p,q)-linear torus curve of a maximal contact. Indeed, $f(x,y)$ takes the form:

$$\begin{aligned} f(x,y) &= \prod_{j=1}^q \left(\zeta^{j-1}y^p - y + x^p\right) \\ &= (y^p)^q - (y-x^p)^q \\ &= (y^q)^p - (y-x^p)^q. \end{aligned}$$

This expression shows that C is a (p,q)-linear torus curve of a maximal contact. □

For practical computations, we suppose hereafter that $\alpha_1,\dots,\alpha_q$ are real numbers such that $\alpha_1 > \cdots > \alpha_q > 0$. Let γ_j be a real positive number such that $\gamma_j^{p-1} = 1/\alpha_j$ for $j=1,\dots,q$. By the assumption $\alpha_1 > \cdots > \alpha_q > 0$, we have

$$0 < \gamma_1 < \cdots < \gamma_q.$$

As $C_j \cap C_i = \{O\}$ for any $j \neq i$, the possible singular pencil $L_s = \{y = s\}$ is either $\{y=0\}$ or L_s is tangent to one of C_j outside of O.

Lemma 2. *Under the above situation, the local data of C for the calculation of the fundamental group of $\mathbb{P}^2 \setminus C$ is the following.*

1. *Singular pencil lines are $y=0$ and $y=\gamma_j\xi^k$ for $j=1,\dots,q$ and $k=0,\dots,p-2$. The pencil lines $y=\gamma_j\xi^k$ is tangent to C_j at $Q_{j,k} := (0,\gamma_j\xi^k)$.*
2. *Two curves C_j and C_i $(j\neq i)$ intersect only at $O \in \mathbb{C}^2$ and $I(C_j,C_i;O) = p^2$.*
3. *The singularity type C at O is given by $(C,O) \sim B_{p^2q,q}$.*

3.2. **Calculation of the fundamental group $\pi_1(\mathbb{P}^2 \setminus C)$ and $\pi_1(\mathbb{C}^2 \setminus C)$.** For the calculations of the fundamental groups $\pi_1(\mathbb{P}^2 \setminus C)$ and $\pi_1(\mathbb{C}^2 \setminus C)$, we use the van Kampen-Zariski pencil method. We take the base point $B_0 = [1:0:0]$ in $\mathbb{P}^2$ and consider the pencil $\mathcal{L} = \{L_s \mid s \in \mathbb{C}\}$ through B_0 with $L_s = \{Y = sZ\}$. The line at infinity L_∞ is given by $\{Z=0\}$. Then L_∞ is generic with respect to C. Affine pencil is $\mathcal{L}^a = \{L_s^a\}_{s\in\mathbb{C}}$ with $L_s^a = \{y=s\}$. (By abuse of notation, we consider this pencil $\mathcal{L} = \{L_s \mid s \in \mathbb{C}\}$ in $\mathbb{C}_t^2$ and $\mathbb{C}_s^2$.) By Lemma 2, the set of parameters $\Sigma \subset \mathbb{C}_y$ which corresponds to singular pencil lines for C is given as follows:

$$\Sigma := \{0, \gamma_j\xi^k \in \mathbb{C}_y \mid k=0,\dots,p-2,\ j=1,\dots,q\}.$$

Take the base point γ_0 of $\mathbb{C}_y \setminus \Sigma$ on the real axis so that $0 < \gamma_0 < \gamma_1$. As

$$\psi(\gamma_0,\alpha_j) - \psi(\gamma_0,\alpha_i) = (\alpha_i - \alpha_j)\gamma_0^p > 0 \quad \text{if} \quad i<j,$$

we have

$$0 < \psi(\gamma_0,\alpha_1) < \psi(\gamma_0,\alpha_2) < \cdots < \psi(\gamma_0,\alpha_q).$$

We take the base point $*_0 = (\tau_0,\gamma_0)$ where τ_0 is a sufficiently large positive number. As C is the pull-back of D by the p-fold cyclic covering $\varphi_p : (x,y) \mapsto (x^p,y)$, the monodromy relations for $\pi_1(L_{\gamma_0}^a \setminus L_{\gamma_0}^a \cap C; *_0)$ are essentially obtained by taking lifting the monodromy relations for $\pi_1(L_{\gamma_0}^a \setminus (L_{\gamma_0}^a \cap D) \cup \{0\}; *_t)$ by φ_p where the base point $*_t$ is a real point defined by $*_t = (\tau_0^p, \gamma_0)$ ([5]). This is the basic idea for the computation of the fundamental groups.

We first take loops $b_1,\dots,b_q$ of $\pi_1(L_{\gamma_0}^a \setminus (L_{\gamma_0}^a \cap D) \cup \{0\}, *_t)$ and put $\tau := b_1\cdots b_q$ as in Figure 2.



FIGURE 2. The loops $b_1,\dots,b_q$ in $\{y=\gamma_0\} \cap \mathbb{C}_t^2$

Let $a'_{i,j}$ and ω'_i be the pull-back of b_j and τ by φ_p respectively starting from $*_i := (\eta^i\tau_0, \gamma_0)$ with $i = 0, \dots, p-1$ where $\eta := \exp(2\pi\sqrt{-1}/p)$ and let $a_{i,j}$ and ω_i be the loop $\ell_i \circ a'_{i,j} \circ \ell_i^{-1}$ and $\ell_i \circ \omega'_i \circ \ell_i^{-1}$ where ℓ_i is the arc of the circle $|x| = \tau_0$ from $*_0$ to $*_i$ as in Figure 3. Hereafter we identify $a'_{i,j}$ and $a_{i,j}$ in this way.



FIGURE 3. The loops $a_{i,j}$ and ω_i in $\{y = \gamma_0\} \cap \mathbb{C}^2_s$

First we see that the monodromy relations on the real axis in $\mathbb{C}_y$ which correspond to singular pencil lines $y = 0$ and $y = \gamma_j$ for $j = 1, \dots, q$. To see these monodromy relations, we consider following loops σ_0 and σ_j in $\mathbb{C}_y$ for $j = 1, \dots, q$. First we define the loop σ_0. Let K_0 be the line segment from γ_0 to $0 - \varepsilon$ on the real axis and let S_0 be the circle $|y| = \varepsilon$ where the circle is always oriented counterclockwise. Then σ_0 is defined as the loop (see Figure 4)

$$\sigma_0 := K_0 \circ S_0 \circ K_0^{-1}.$$

Next we define loops σ_j for $j = 1, \dots, q$. Let S_j be the loop which is represented by the circle $|y - \gamma_j| = \varepsilon$ oriented counter clockwise. Let K_j be the modified line segment from γ_0 to $\gamma_j - \varepsilon$. The segment $[\gamma_i - \varepsilon, \gamma_i + \varepsilon]$ is replaced by the lower half circle of S_i. Then σ_j is defined as the loop (see Figure 4)

$$\sigma_j := K_j \circ S_j \circ K_j^{-1}.$$



FIGURE 4. Loops in $\mathbb{C}_y$

Case 1: First we see the monodromy relations at $y = 0$. By the definitions of C_j's and Lemma 2, the origin O is a flex point of C_j such that $\{y = 0\}$ is the tangent line for $j = 1, \dots, q$ and C_i and C_j intersect with intersection multiplicity p^2 at O for each $i \neq j$ and the topological type of C at O is $B_{p^2q,q}$.

To see that monodromy relations, we look at the Puiseux parametrization of each component C_j at O. Consider that curves D_j and D whose defining polynomials are $g_j(x,y) = x - \psi(y,\alpha_j)$ and $g(x,y) = \prod_{j=1}^{q} g_j(x,y)$ respectively. By the definitions, $\psi(y,\alpha_j) = y(y - \alpha_j y^{p-1})$, $f_j(x,y) = g_j(x^p,y)$, we have $x^p = y(1-\alpha_j y^{p-1})$. By the generalized binomial theorem, we can solve $x^p = y(1-\alpha_j y^{p-1})$ as follows.

$$(1)\quad C_j : \quad \begin{cases} x = \varphi_j(t), \quad \varphi_j(t) = t\left(1 - \dfrac{\alpha_j}{p}\, t^{p(p-1)} + \cdots\right), & j = 1,\dots,q. \\ y = t^p, \end{cases}$$

$$(2)\quad \frac{\varphi_j(t)}{t} - \frac{\varphi_i(t)}{t} = \frac{1}{p}(\alpha_i - \alpha_j)t^{p(p-1)} + \cdots, \quad j \neq i.$$

Note that the leading term of $\varphi_j(t)$ is t which is independent of index $j = 1,\dots,q$. The topological behavior of the centers of the generators, pq points $C \cap \{y = \varepsilon\exp(\sqrt{-1}\,\theta)\}$, looks like the movements of satellites around planets with $0 \le \theta \le 2\pi$. For a fixed y, there are p choices of t so that $y = t^p$. We take t so that $0 \le \arg t \le 2\pi/p$. Thus planets are the points $P_i = (t\eta^i, t^p)$ for $i = 0,\dots,p-1$ and the satellites around P_i are $\{(\varphi_j(t\eta^i), t^p) \,|\, j = 1,\dots,q\}$ where $\eta = \exp(2\pi\sqrt{-1}/p)$.

Above conditions (1) and (2) say that p planets moves an arc of the angle $2\pi/p$ centered at the origin when $t = \varepsilon^{1/p}\exp(\sqrt{-1}\,\theta/p)$ moves from $\theta = 0$ to 2π. Then the satellites, which are the center of loops $\{a_{i,j} \,|\, j = 1,\dots,q\}$, are rotated $(p-1)$-times around P_i simultaneously for $i = 0,\dots,p-1$. Hence we have the monodromy relations:

$$(1\text{-}1)\quad a_{i,j} = a_{i,j}^{\sigma_0} = \begin{cases} \omega_{i+1}^{p-1} a_{i+1,j} \omega_{i+1}^{-(p-1)} & 0 \le i \le p-2, \\ \Omega\,\omega_1^{p-1} a_{0,j} (\Omega\,\omega_1^{p-1})^{-1} & i = p-1, \end{cases} \qquad j = 1\dots,q$$

where $a_{i,j}^{\sigma_0}$ is the monodromy action by σ_0 on $a_{i,j}$. See Figure 5 for the case $p = 3$ and $q = 2$.



FIGURE 5. The case $p = 3$ and $q = 2$

On the other hand, we get the relation $\omega_1 = \omega_2 = \cdots = \omega_p$ when $y = \varepsilon\exp(2\pi\sqrt{-1}\theta)$ moves around the origin once. Hence we have

$$\Omega = \omega^p, \quad \omega := \omega_1 = \omega_2 = \cdots = \omega_p.$$

We can rewrite the relations (1-1) as follows:

$$(1\text{-}2)\quad a_{i,j} = \begin{cases} \omega^{p-1} a_{i+1,j} \omega^{-(p-1)} & 0 \le i \le p-2, \\ \omega^{2p-1} a_{0,j} \omega^{-(2p-1)} & i = p-1, \end{cases} \qquad j = 1,\dots,q.$$

Case 2: Next we consider the monodromy relations at $y = \gamma_j$ for $j \geq 1$. In this case, the pencil line L_{γ_j} is tangent to C_j and $C_j \cap L_{\gamma_j} = \{Q_{j,0}\} = \{(0, \gamma_j)\}$ is a flex point of C_j of flex order $p-2$. On the other hand, the pencil line L_{γ_j} is generic with respect to other C_i for $i \neq j$.

First we consider the case $j = 1$. Recall that the defining polynomial of C_i is

$$f_i(x,y) = x^p - \psi(y, \alpha_i) = x^p - y\prod_{k=0}^{p-2}(\gamma_i \xi^k - y), \quad i = 1, \dots, q.$$

We take the local coordinates $(x, y_1) := (x, y - \gamma_1)$ centered at $Q_{1,0}$. By an easy calculation,

$$f_i(x, y_1 + \gamma_1) = 0 \Leftrightarrow x^p = \begin{cases} (1-p)y_1 + H_1(y_1), & i = 1, \\ \frac{\gamma_1}{\alpha_1}(\alpha_1 - \alpha_i) + H_i(y_1) & i \neq 1 \end{cases} \tag{1}$$

and $\alpha_1 - \alpha_i > 0$ where $\operatorname{ord}_{y_1} H_1 \geq 2$ and $\operatorname{ord}_{y_1} H_i \geq 1$ for $i \geq 2$. The first coefficients $1-p$ and $\frac{\gamma_1}{\alpha_1}(\alpha_1 - \alpha_i)$ are obtained from the equalities:

$$\begin{cases} \psi(y_1 + \gamma_1, \alpha_1) = -(y_1 + \gamma_1)y_1 \prod_{k \geq 1}(\gamma_1 \xi^k - y_1 - \gamma_1) \\ \psi(y_1 + \gamma_1, \alpha_i) = (y_1 + \gamma_1) - \alpha_i (y_1 + \gamma_1)^p \end{cases}$$

and

$$\begin{cases} \frac{d\psi(y_1+\gamma_1, \alpha_1)}{dy_1}|_{y_1=0} = \frac{d\psi(y, \alpha_1)}{dy}|_{y=\gamma_1} = 1-p \\ \psi(\gamma_1, \alpha_i) = \frac{\gamma_1}{\alpha_1}(\alpha_1 - \alpha_i). \end{cases}$$

Now we consider the monodromy relations at $y = \gamma_1$. First, the action of σ_1 on $b_1, \dots, b_q$ is sketched as in Figure 6. Thus we see that the generators which are topologically deformed are $\{a_{i,1} \mid i = 0, \dots, p-1\}$ under the rotation $y_1 = -\varepsilon \exp(\sqrt{-1}\theta)$ with $0 \leq \theta \leq 2\pi$. The other generators are unchanged. Namely $a_{i,j}^{\sigma_1} = a_{i,j}$ for $i = 0, \dots, p-1$ and $j \geq 2$. To simplify the monodromy relations, we introduce an element $g_1 := b_2 \cdots b_q$. Then $\tau = b_1 g_1$. See Figure 6.



FIGURE 6. $\{y = \gamma_1 - \varepsilon\} \cap \mathbb{C}_t^2$

Let $g_{i,1}$ be the pull-back of g_1 starting from $*_i$ for $i = 0, \dots, p-1$. More precisely, $g_{i,1} = a_{i,2} \cdots a_{i,q}$ and $\omega_i = a_{i,1} g_{i,1}$.
When $y_1 = y - \gamma_1 = -\varepsilon \exp(\sqrt{-1}\theta)$ moves from $\theta = 0$ to 2π, the generators $a_{0,1}, \dots, a_{p-1,1}$ moves an arc of the angle $2\pi/p$ centered at the origin (the lifts of $b_1^{\sigma_1}$) and the other generators do not move. Thus we have following monodromy relations:

$$a_{i,1} = a_{i,1}^{\sigma_1} = \begin{cases} g_{i+1,1}^{-1}\, a_{i+1,1}\, g_{i+1,1} & 0 \leq i \leq p-2 \\ \Omega\, g_{0,1}^{-1}\, a_{0,1}\, (\Omega\, g_{0,1}^{-1})^{-1} & i = p-1 \end{cases} \tag{2-1}$$

and $g_{i,1}^{\sigma_1} = g_{i,1}$ for $i = 0, \dots, p-1$. See Figure 7.



FIGURE 7. The action of σ_1

By the previous argument, we have $\omega_1 = \omega$ and $\Omega = \omega^p$. Hence we can rewrite the relations (2-1) as follows:

$$a_{i,1} = a_{i,1}^{\sigma_1} = \begin{cases} \omega^{-1} a_{i+1,1} \omega & 0 \le i \le p-2 \\ \omega^{p-1} a_{0,1} \omega^{-(p-1)} & i = p-1 \end{cases} \tag{2-2}$$

Now we consider the case $j \ge 2$. First we deform the pencil from γ_0 to $\gamma_j - \varepsilon$ along K_j. Note that

$$\psi(y, \alpha_k) \begin{cases} < 0 & k < j \\ > 0 & k > j. \end{cases}$$

where $y \in [\gamma_{j-1} + \varepsilon, \gamma_j - \varepsilon]$. Thus the generators $b_1, \dots, b_q$ are deformed as in Figure 8 where $\psi_j := \psi(y, \alpha_j)$.



FIGURE 8. $\{y = \gamma_j - \varepsilon\} \cap \mathbb{C}_t^2$

When y moves along $S_j : |y - \gamma_j| = \varepsilon$, the single root of $g_{\alpha_j}(x, y) = 0$ which is near the origin goes around the origin once and the other roots $g_{\alpha_k}(x, y) = 0$ $(k \neq j)$ do not move as in Figure 9 where $\psi_j := \psi(y, \alpha_j)$.



FIGURE 9. $\{y = \gamma_j - \varepsilon\} \cap \mathbb{C}_t^2$

This implies, by taking p-fold covering, the corresponding generators $a_{0,j}, \dots, a_{p-1,j}$ of b_j moves an arc of the angle $2\pi/p$ centered at the origin.

To see it more precisely, we put new loops:

$$h_j = \begin{cases} e & j = 1 \\ b_1 \cdots b_{j-1} & 2 \le j \le q \end{cases}, \qquad g_j = \begin{cases} b_{j+1} \cdots b_q & 1 \le j \le q-1 \\ e & j = q. \end{cases}$$

By the definitions, we have $\tau = h_j b_j g_j$. See Figure 10.



FIGURE 10. New loops

We take the local coordinates $(x, y_j) := (x, y - \gamma_j)$ centered at $Q_{j,0}$. Then

$$f_i(x, y_j + \gamma_j) = 0 \Leftrightarrow x^p = \begin{cases} (1-p)y_j + H_j(y_j), & i = j, \\ \frac{\gamma_j}{\alpha_j}(\alpha_j - \alpha_i) + H_i(y_j) & i \neq j \end{cases}$$

where $\operatorname{ord}_{y_j} H_j \geq 2$ and $\operatorname{ord}_{y_j} H_i \geq 1$ for $i \neq j$. By the assumption, we have $\alpha_j - \alpha_i > 0$ or $\alpha_j - \alpha_i < 0$ corresponding to either $i > j$ or $i < j$ respectively. Thus we can see that the generators which are deformed under this monodromy are $\{a_{i,j} \mid i = 0, \dots, p-1\}$ when y_j moves around the circle $|y_j| = \varepsilon$. Thus $a_{i,k}^{\sigma_j} = a_{i,k}$ for $k \neq j$.
Let $h_{i,j}$ and $g_{i,j}$ be the pull-back of h_j and g_j respectively. By the definition, we have

$$h_{i,j} = a_{i,1} \cdots a_{i,j-1}$$
$$g_{i,j} = a_{i,j+1} \cdots a_{i,q}$$

where $h_{i,1} = e$ and $g_{i,q} = e$ and we put $\omega_i = h_{i,j} a_{i,j} g_{i,j}$.

When y moves around the circle $|y - \gamma_j| = \varepsilon$ once, the generators $a_{0,j}, \dots, a_{p-1,j}$ moves an arc of the angle $2\pi/p$ centered at the origin. Thus we have following monodromy relations:

$$(2\text{-}3) \quad a_{i,j} = a_{i,j}^{\sigma_j} = \begin{cases} (g_{i+1,j} h_{i,j})^{-1}\, a_{i+1,j}\, g_{i+1,j} h_{i,j} & 0 \leq i \leq p-2 \\ h_{p-1,j}^{-1}\, \Omega\, g_{0,j}^{-1}\, a_{0,j}\, (h_{p-1,j}^{-1} \Omega g_{0,j}^{-1})^{-1} & i = p-1 \end{cases}$$

and $g_{i,j}^{\sigma_j} = g_{i,j}$ and $h_{i,j}^{\sigma_j} = h_{i,j}$ for $i = 0, \dots, p-1$. See Figure 11.



FIGURE 11. The action of σ_j

Other cases: Finally we read the monodromy relations at $y = \gamma$ where $\gamma \in \Sigma$ with $\gamma \neq 0, \gamma_1, \dots, \gamma_q$. Recall that the set of parameters $\Sigma \subset \mathbb{C}_y$ which corresponds to singular pencils are given by

$$\Sigma = \{0, \gamma_j \xi^k \in \mathbb{C}_y \,|\, k = 0, \dots, p-2,\ j = 1, \dots, q\}, \quad \xi = \exp\left(\frac{2\pi\sqrt{-1}}{p-1}\right).$$

Then the pencil line $L_{\gamma_j \xi^k} = \{y = \gamma_j \xi^k\}$ is singular with respect to C_j and $C_j \cap L_{\gamma_j \xi^k} = \{Q_{j,k}\} = \{(0, \gamma_j \xi^k)\}$ is a flex point of C_j of flex order $p-2$ for $k = 1, \dots, p-2$. Note that the pencil line $L_{\gamma_j \xi^k}$ is generic with respect to other C_i for $i \neq j$.

First we consider the case $k = 1$. That is, we consider the monodromy relations at $y = \gamma_j \xi$. We take a path L_1 which connects γ_0 and $\gamma_0 \xi$ as in Figure 12.



FIGURE 12. The loop L_1

Then the loops $b_1, \dots, b_q$ are deformed as in the left side of Figure 13.We take new loops $c_1, \dots, c_q$ as in the right side of Figure 13. Here $\xi *_t = (\xi \omega_0^p, \xi \gamma_0)$.



FIGURE 13. New loops $c_1, \dots, c_q$.

They are related by the following.

$$c_j = \tau^{-1} b_j \tau, \quad j = 1, \dots, q. \tag{2}$$

Let $d_{0,j}, \dots, d_{p-1,j}$ be the pull-back of c_j by φ_p for $j = 1, \dots, q$. Then the relation (2) implies

$$d_{i,j} = \omega^{-1} a_{i,j} \omega. \tag{3}$$

Now we consider the loops $\sigma_1^{(1)}, \dots, \sigma_q^{(1)}$ in $\mathbb{C}_y$ with base point $\gamma_0 \xi$ as in Figure 14.



FIGURE 14. The loop $\sigma_j^{(1)}$

We will see that the monodromy relations are exactly as (2-3). To see this assertion, we take the following modified coordinates $(\tilde{x}, \tilde{y})$ which are defined by

$$\tilde{x} := \exp\left(\frac{-2\pi\sqrt{-1}}{p(p-1)}\right)x, \quad \tilde{y} := \xi y.$$

In these coordinates, the loops $\sigma_1^{(1)}, \dots, \sigma_q^{(1)}$ coincide with $\sigma_1, \dots, \sigma_q$ and C_j is defined by the same equality:

$$C_j : \quad \tilde{x}^p = \tilde{y}(1 - \alpha_j \tilde{y}^{p-1}).$$

The situation of loops $c_1, \dots, c_q$ are the same with that of $b_1, \dots, b_q$ and the situation of loops $d_{i,j}, i = 0, \dots, p-1, j = 1, \dots, q$ are the same with that of $a_{i,j}, i = 0, \dots, p-1, j = 1, \dots, q$. Therefore we obtain the relations

$$(2\text{-}3)' \quad d_{i,j} = \begin{cases} (\tilde{g}_{i+1,j}\tilde{h}_{i,j})^{-1}\, d_{i+1,j}\, \tilde{g}_{i+1,j}\tilde{h}_{i,j} & 0 \le i \le p-2 \\ \tilde{h}_{p-1,j}^{-1}\, \tilde{\Omega}\, \tilde{g}_{0,j}^{-1}\, d_{0,j}\, (\tilde{h}_{p-1,j}^{-1}\tilde{\Omega}\tilde{g}_{0,j}^{-1})^{-1} & i = p-1 \end{cases}, \quad j = 1, \dots, q$$

where $\tilde{h}_{i,j} := d_{i,1} \cdots d_{i,j-1}$, $\tilde{g}_{i,j} := d_{i,j+1} \cdots d_{i,q}$ and $\tilde{\Omega} := \omega^{-1}\Omega\omega$. Now we claim the following.

Lemma 3. *The relation* (2-3)′ *is the same with the relation* (2-3).

Proof. First we consider the relation $d_{i,j} = (\tilde{g}_{i+1,j}\tilde{h}_{i,j})^{-1}\, d_{i+1,j}\, \tilde{g}_{i+1,j}\tilde{h}_{i,j}$ in (2-3)′. By the relation (3), we have

$$\tilde{h}_{i,j} = \omega^{-1}h_{i,j}\omega, \quad \tilde{g}_{i,j} = \omega^{-1}g_{i,j}\omega.$$

Thus $d_{i,j} = (\tilde{g}_{i+1,j}\tilde{h}_{i,j})^{-1}\, d_{i+1,j}\, \tilde{g}_{i+1,j}\tilde{h}_{i,j}$ can be translated as follows

$$\begin{aligned} d_{i,j} = \omega^{-1}a_{i,j}\omega &= (\tilde{g}_{i+1,j}\tilde{h}_{i,j})^{-1}\, d_{i+1,j}\, \tilde{g}_{i+1,j}\tilde{h}_{i,j} \\ &= \left((\omega^{-1}g_{i+1,j}\omega)(\omega^{-1}h_{i,j}\omega)\right)^{-1}(\omega^{-1}a_{i+1,j}\omega)(\omega^{-1}g_{i+1,j}\omega)(\omega^{-1}h_{i,j}\omega) \\ &= \omega^{-1}(g_{i+1,j}h_{i,j})^{-1}a_{i+1,j}g_{i+1,j}h_{i,j}\omega \end{aligned}$$

which implies (2-3). For the relation $d_{i,j} = \tilde{h}_{p-1,j}^{-1}\, \tilde{\Omega}\, \tilde{g}_{0,j}^{-1}\, d_{0,j}\, (\tilde{h}_{p-1,j}^{-1}\tilde{\Omega}\tilde{g}_{0,j}^{-1})^{-1}$, the argument is the same. This complete the proof. □

Next we consider general cases $k \ge 2$. That is, we consider the monodromy relations at $y = \gamma_j \xi^k$. Then we take a path L_k which connects γ_0 and $\gamma_0\xi^k$:



FIGURE 15. The loop L_k

By the exact same arguments as in the case $k = 1$, we see that no new monodromy relations are necessary.

3.3. The group structures of $\pi_1(\mathbb{C}^2 \setminus C)$ and $\pi_1(\mathbb{P}^2 \setminus C)$.

In this section, we consider the group structures of $\pi_1(\mathbb{P}^2 - C)$ and $\pi_1(\mathbb{C}^2 \setminus C)$. First by previous considerations, we have prove that

$$\pi_1(\mathbb{C}^2 \setminus C) = \langle \omega,\, a_{i,j}, i = 0, \dots, p-1, j = 1, \dots, q, \mid \text{(1-2)}, \text{(2-3)}, (S) \rangle \tag{4}$$

where

$$\text{(1-2)} \quad a_{i,j} = \begin{cases} \omega^{p-1} a_{i+1,j} \omega^{-(p-1)} & 0 \le i \le p-2, \\ \omega^{2p-1} a_{0,j} (\omega^{2p-1})^{-1} & i = p-1, \end{cases}, \quad j = 1, \dots, q$$

$$\text{(2-3)} \quad a_{i,j} = \begin{cases} (g_{i+1,j} h_{i,j})^{-1}\, a_{i+1,j}\, g_{i+1,j} h_{i,j} & 0 \le i \le p-2 \\ h_{p-1,j}^{-1}\, \Omega\, g_{0,j}^{-1}\, a_{0,j}\, (h_{p-1,j}^{-1} \Omega g_{0,j}^{-1})^{-1} & i = p-1 \end{cases}, \quad j = 1, \dots, q$$

$$(S) \qquad \omega = a_{0,1} \cdots a_{0,q}.$$

Note that last relations in (1-2) and (2-3) are unnecessary as they follow from previous relations. By the definitions of $g_{i,j}$ and $h_{i,j}$ and (1-2), we have the following inductive relations.

$$\begin{cases} h_{i+1,j} = \omega^{-(p-1)} h_{i,j} \omega^{p-1}, \quad g_{i+1,j} = \omega^{-(p-1)} g_{i,j} \omega^{p-1} \\ g_{i+1,j} h_{i,j} = \omega^{-(p-1)} g_{i,j} \omega^{p-1} h_{i,j} \end{cases} \tag{5}$$

First we examine the relation (2-3) for a fixed $i \le p-2$ using (1-2). The case $j = 1$ gives the equality:

$$\begin{aligned} a_{i,1} &= (g_{i+1,1})^{-1}\, a_{i+1,1}\, g_{i+1,1}, \qquad \text{as } h_{i,1} = e \\ &= (\omega^{-(p-1)} g_{i,1} \omega^{p-1})^{-1} (\omega^{-(p-1)} a_{i,1} \omega^{p-1})(\omega^{-(p-1)} g_{i,1} \omega^{p-1}) \\ &= \omega^{-(p-1)} g_{i,1}^{-1} a_{i,1} g_{i,1} \omega^{p-1} \\ &= \omega^{-p} a_{i,1} \omega^{p}. \end{aligned}$$

This implies ω^p and $a_{i,1}$ commute. Now by the induction on j, we show that

$$(R_i) \qquad [a_{i,j}, \omega^p] = e, \qquad j = 1, \dots, q$$

where $[a, b] = aba^{-1}b^{-1}$.

Proof. In fact, assuming $a_{i,1}, \dots, a_{i,j-1}$ commute with ω^p, we get

$$\begin{aligned} a_{i,j} &= (g_{i+1,j} h_{i,j})^{-1}\, a_{i+1,j}\, g_{i+1,j} h_{i,j} \\ &= (\omega^{-(p-1)} g_{i,j} \omega^{p-1} h_{i,j})^{-1} (\omega^{-(p-1)} a_{i,j} \omega^{p-1})(\omega^{-(p-1)} g_{i,j} \omega^{p-1} h_{i,j}) \\ &= h_{i,j}^{-1} \omega^{-(p-1)} g_{i,j}^{-1} a_{i,j} g_{i,j} \omega^{p-1} h_{i,j} \\ &= h_{i,j}^{-1} \omega^{-(p-1)} g_{i,j}^{-1} (h_{i,j}^{-1} h_{i,j}) a_{i,j} g_{i,j} \omega^{p-1} h_{i,j} \\ &= h_{i,j}^{-1} \omega^{-p} h_{i,j} a_{i,j} h_{i,j}^{-1} \omega^{p} h_{i,j}, \qquad \text{as } [h_{i,j}, \omega^p] = e \\ &= \omega^{-p} a_{i,j} \omega^{p}. \end{aligned}$$

Thus we get $[a_{i,j}, \omega^p] = e$ for all $j = 1, \dots, q$. □

The relation (R_i) for $i = 0, \dots, p-1$ implies ω^p is in the center of $\pi_1(\mathbb{C}^2 \setminus C)$. Using relations (1-2) and (R_i), we have

$$a_{i+1,j} = \omega a_{i,j} \omega^{-1}, \quad i = 0, \dots, p-2, \quad j = 1, \dots, q.$$

Thus we get

$$a_{i,j} = \omega^i a_{0,j} \omega^{-i}, \quad i = 0, \dots, p-1, \quad j = 1, \dots, q. \tag{6}$$

Hence we can take $a_{0,1}, \dots, a_{0,q}$ as generators. They satisfy the relations

$$[a_{0,j}, \omega^p] = e, \quad j = 1, \dots, q. \tag{R_0}$$

It is easy to see (and we have seen implicitly in the above discussions) that the relations (1-2) and (2-3) follow from (R_0), (S) and (5). Thus we have shown

$$\begin{aligned}
\pi_1(\mathbb{C}^2 \setminus C) &= \langle a_{i,j}\,(i = 0, \dots, p-1, j = 1, \dots, q), \omega \,|\, \text{(1-1)},\ \text{(2-3)},\ (S),\ (5) \rangle \\
&\cong \langle a_{0,1}, \dots, a_{0,q}, \omega \,|\, (R_0),\ (S) \rangle \\
\pi_1(\mathbb{P}^2 \setminus C) &\cong \langle a_{0,1}, \dots, a_{0,q}, \omega \,|\, \omega^p = e,\ (R_0),\ (S) \rangle \\
&\cong \langle a_{0,1}, \dots, a_{0,q}, \omega \,|\, \omega^p = e,\ (S) \rangle \\
&\cong \langle a_{0,1}, \dots, a_{0,q-1}, \omega \,|\, \omega^p = e \rangle \\
&\cong F(q-1) * \mathbb{Z}/p\mathbb{Z}.
\end{aligned}$$

This completes the proof of Theorem 1.

I would like to express my deepest gratitude to Professor Mutsuo Oka for suggesting me the present problem and helping me to prepare this paper.

REFERENCES

[1] B. Audoubert, C. Nguyen, and M. Oka. On alexander polynomials of torus curves. *J. Math. Soc. Japan*, 57(4):935–957, 2005.

[2] B. Audoubert, T. C. Nguyen, and M. Oka. Erratum to: "On Alexander polynomials of torus curves" [J. Math. Soc. Japan **57** (2005), no. 4, 935–957; mr2183580]. *J. Math. Soc. Japan*, 59(3):911–912, 2007.

[3] M. Oka. On the fundamental group of the complement of a reducible curve in P^2. *J. London Math. Soc. (2)*, 12(2):239–252, 1975/76.

[4] M. Oka. On the fundamental group of the complement of certain plane curves. *J. Math. Soc. Japan*, 30(4):579–597, 1978.

[5] M. Oka. Two transforms of plane curves and their fundamental groups. *J. Math. Sci. Univ. Tokyo*, 3:399–443, 1996.

[6] M. Oka. A survey on Alexander polynomials of plane curves. *Singularités Franco-Japonaise, Séminaire et congrès*, 10:209–232, 2005.

[7] E. van Kampen. On the fundamental group of an algebraic curve. *Amer. J. Math.*, 55:255–260, 1933.

[8] O. Zariski. On the poincaré group of a rational plane curves. *Amer. J. Math.*, 58:607–619, 1929.

[9] O. Zariski. On the problem of existence of algebraic functions of two variables possessing a given branch curve. *Amer. J. Math.*, 51:305–328, 1929.

DEPARTMENT OF MATHEMATICS
TOKYO UNIVERSITY OF SCIENCE
WAKAMIYA-CHO 26, SHINJUKU-KU
TOKYO 162-0827 JAPAN
E-mail: j1107702@ed.kagu.tus.ac.jp